ELSEVIER

Environmental Pollution 117 (2002) I–V

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 116

Number 1 2001

Editorial

1 A new cover: a new view of the world
W.J. Manning

3 Distribution of ozone and other air pollutants in forests of the Carpathian Mountains in central Europe
A. BYTNEROWICZ (USA), B. GODZIK (Poland), W. FRACZEK (USA), K. GRODZIŃSKA, M. KRYWULT (Poland), O. BADEA (Romania), P. BARANČOK (Slovakia), O. BLUM (Ukraine), M. ČERNY (Czech Republic), S. GODZIK (Poland), B. MANKOVSKA (Slovakia), W. MANNING (USA), P. MORAVČIK (Czech Republic), R. MUSSELMAN (USA), J. OSZLANYI (Slovakia), D. POSTELNICU (Romania), J. SZDŹUJ (Poland), M. VARŠAVOVA (Slovakia), M. ZOTA (Romania)
Models of ozone distribution and evaluation of its phytotoxic potential in the forests of the Carpathian Mountains of Central Europe are presented.

27 Ozone impacts on loblolly pine (*Pinus taeda* L.) grown in a competitive environment
D.N. BARBO, A.H. CHAPPELKA, G.L. SOMERS, M.S. MILLER-GOODMAN, K. STOLTE (USA)
Loblolly pine seedlings in carbon-filtered air open-top chambers, with competition from other plants, behaved differently than those reported in other studies without competition.

37 Impact of fungicides on active oxygen species and antioxidant enzymes in spring barley (*Hordeum vulgare* L.) exposed to ozone
Y.-X. WU, A. VON TIEDEMANN (Germany)
Modern fungicides widely used in cereal production have antioxidant effects and may reduce the impact of ozone on grain production.

49 Distribution of organochlorine pesticides in seawater of the Bering and Chukchi Sea
Y. ZI-WEI, J. GUI-BIN, X. HENG-ZHEN (People's Republic of China)
Organochlorine pesticides were determined, with many different residue patterns found for the first time in this region.

57 A fragment constant QSAR model for evaluating the EC_{50} values of organic chemicals to *Daphnia Magna*
S. TAO, X. XI, F. XU, B. LI, J. CAO, R. DAWSON (China)
Effects of toxic chemicals can be estimated from quantitative struture–activity relationships (QSARs).

65 Spatial distribution of mercury in the sediments and riparian environment of the River Yare, Norfolk, UK
J.W. BIRKETT, J.M.K. NORENG, J.N. LESTER (UK)
The spatial distribution of mercury follows a distinct pollution plume downstream of a sewage outfall.

75 Net fluxes of pesticides from the Scheldt Estuary into the North Sea: a model approach
R.J.C.A. STEEN, E.H.G. EVERS, B. VAN HATTUM, W.P. COFINO, U.A.T. BRINKMAN (The Netherlands)
Pesticide fluxes are modeled hydrodynamically for a complex estuarine environment.

85 Examination of the bioaccumulation of halogenated dimethyl bipyrroles in an Arctic marine food web using stable nitrogen isotope analysis
S.A. TITTLEMIER, A.T. FISK, K.A. HOBSON, R.J. NORSTROM (Canada)
Except for ringed seals, halogenated dimethyl bipyrrole (HDBP) congeners biomagnified in a sample aquatic food web of invertebrate > fish > seabird.

95 Parallelisation and application of a multi-layer atmospheric transport model to quantify dispersion and deposition of ammonia over the British Isles
N. FOURNIER (UK), V.A. PAIS (Italy), M.A. SUTTON, K.J. WESTON, U. DRAGOSITS, S.Y. TANG (UK), J. AHERNE (Ireland)
A parallel version of an atmospheric transport model (FRAME) is developed and applied to the British Isles for the first time.

109 Potentials and drawbacks of chelate-enhanced phytoremediation of soils
P. RÖMKENS, L. BOUWMAN, J. JAPENGA, C. DRAAISMA (The Netherlands)
This paper reviews chelate-enhanced phytoremediation of heavy metals, focusing on unwanted side-effects.

123 Biodiversity of epiphytic lichens and air pollution in the town of Siena (Central Italy)
S. LOPPI (Italy), D. IVANOV (Bulgaria), R. BOCCARDI (Italy)
Biodiversity of epiphytic lichens was used to demonstrate improvement of air quality in Siena, Italy.

129 Element concentrations in the forest moss *Hylocomium splendens*: variation associated with altitude, net primary production and soil chemistry
R. GERDOL, L. BRAGAZZA, R. MARCHESINI (Italy)
Metal concentrations in forest mosses may be influenced by variation in net primary production associated with altitude

137 Biological responses to the reversal of acidification in surface waters of the English Lake District
E. TIPPING, J.A.B. BASS, D. HARDIE, E.Y. HAWORTH, M.A. HURLEY, G. WILLS (UK)
Changes in macroinvertebrate and diatom communities are consistent with increases in surface water pH.

PII: S0269-7491(01)00327-X

147 Metal-induced tolerance in the freshwater microbenthic diatom *Gomphonema parvulum*
N. IVORRA, C. BARRANGUET, M. JONKER, M.H.S. KRAAK, W. ADMIRAAL (The Netherlands)
The benthic diatom Gomphonema parvulum *developed zinc tolerance that persisted 2 years after exposure to zinc.*

159 Metal bioavailability and toxicity through a sediment core
U. BORGMANN, W.P. NORWOOD (Canada)
Metal bioaccumulation, relative bioavailability, and toxicity to Hyalella azteca *varied with depth and age of deposition of sediment core sections.*

169 Environmental contaminants in prey and tissues of the peregrine falcon in the Big Bend Region, Texas, USA
M. MORA, R. SKILES, B. MCKINNEY, M. PAREDES, D. BUCKLER, D. PAPOULIAS, D. KLEIN (USA)
Falcon productivity rates in Big Bend may be affected by mercury and selenium.

177 Ecotoxicological studies of environmental samples from Buenos Aires area using a standardized amphibian embryo toxicity test (AMPHITOX)
J. HERKOVITS, C. PEREZ-COLL, F.D. HERKOVITS (Argentina)
AMPHITOX was found to be useful in evaluating toxicity of a wide variety of samples.

Number 2 2001

185 Accumulation of lead and cadmium in the marine prosobranch *Nerita saxtilis*, chemical analysis, light and electron microscopy
A.T. ABDALLAH, M.A. MOUSTAFA (Egypt)
The marine snail Nerita saxtilis may be a useful biomonitor for heavy metals in the Red Sea.

193 UV Fourier transform measurements of tropospheric O_3, NO_2, SO_2, benzene, and toluene
A.C. VANDAELE, A. TSOULI, M. CARLEER, R. COLIN (Belgium)
UV Fourier transform spectroscopy is a powerful tool to study the tropospheric atmosphere.

203 Nematode community response to fish-farm impact in the western Mediterranean
S. MIRTO, T. LA ROSA, C. GAMBI, R. DANOVARO, A. MAZZOLA (Italy)
Nematode analysis in terms of k-dominance, maturity index and genus composition provides a sensible tool for describing the environmental impact due to fish-farm biodeposition in the Mediterranean.

215 Salt secretion and stomatal behaviour in *Avicennia marina* seedlings fumigated with the volatile fraction of light Arabian crude oil
T. YOUSSEF, A. GHANEM (United Arab Emirates)
Light Arabian crude oil fumes reduced salt secretion and transpiration in mangrave seedlings.

225 Water-soluble aerosol and visibility degradation in Hong Kong during autumn and early winter, 1998
Y.L. LEE, R. SEQUEIRA (Hong Kong)
Fine ammoniated sulfate aerosol is probably the predominant water-soluble, inorganic, visibility-reducing component in Hong Kong.

235 Lichen response to changes in atmospheric sulphur: isotopic evidence
R.D. WISEMAN, M.A. WADLEIGH (Canada)
In 18 months, lichens transplanted from a remote area will resemble those in an urban area in terms of sulphur concentration and isotopic composition.

243 Toxicity of uranium mine receiving waters to early life stage fathead minnows (*Pimephales promelas*) in the laboratory
G.G. PYLE, S.M. SWANSON, D.M. LEHMKUHL (Canada)

257 The influence of meteorological conditions on the behavior of pollutants concentrations in São Paulo, Brazil
O.R. SÁNCHEZ-CCOYLLO, M. DE FÁTIMA ANDRADE (Brazil)

265 Accumulation and tissue distribution of mercury and selenium in striped dolphins (*Stenella coeruleoalba)* from the Mediterranean Sea (southern Italy)
N. CARDELLICCHIO, A. DECATALDO, A. DI LEO, A. MISINO (Italy)

273 Urban polluted forest soils induce elevated root peroxidase activity in Scots pine (*Pinus sylvestris* L.) seedlings
A.M. MARKKOLA, O. TARVAINEN (Finland), U. AHONEN-JONNARTH (Sweden), R. STRÖMMER (Finland)
Peroxidase levels in seedlings planted in urban soils were higher than in non-polluted soils.

279 Mosses and lichens as biomonitors of trace metals. A comparison study on *Hypnum cupressiforme* and *Parmelia caperata* in a former mining district in Italy
R. BARGAGLI, F. MONACI, F. BORGHINI, F. BRAVI, C. AGNORELLI (Italy)
Epiphytic lichens accumulated higher concentrations of atmospheric elements than epigeic mosses in a region (Colline Metalifere) with widespread natural and anthropogenic sources of metals.

289 The effect of ageing on the toxicity of zinc for the potworm *Enchytraeus albidus*
K. LOCK, C.R. JANSSEN (Belgium)
No effect of ageing on toxicity of zinc for the potworm Enchytraeus albidus was detected in artificial soil.

293 Plant accumulation of potentially toxic elements in sewage sludge as affected by soil organic matter level and mycorrhizal fungi
M. OUDEH (Syria), M. KHAN (Pakistan), J. SCULLION (UK)
Mycorrhizal activity and soil organic matter levels have the potential to modify the risks for plants from metals in sludges.

301 Treated wood in livestock facilities: relationships among residues of pentachlorophenol, dioxins, and furans in wood and beef
G.F. FRIES, V.J. FEIL, R.G. ZAYLSKIE, K.M. BIALEK, C.P. RICE (USA)
Ingestion of PCP-treated wood is shown to be a potentially important source of PCDD/F residues in US cattle.

309 Fish community responses to pulp and paper mill effluents at the southern Lake Saimaa, Finland
A.E. KARELS, A. NIEMI (Finland)
Fish responses to pulp and paper mill effluents appear related to habitat changes caused by nutrient enrichment.

319 Organochlorine pesticides and hexachlorobenzene along the central coast of New South Wales: multi-scale distributions using the territorial damselfish *Parma microlepis* as an indicator
K.A. TRICKLEBANK, M.J. KINGSFORD, H.A. ROSE (Australia)
Clear differences in patterns of distribution of organochlorines were found between urban and reference locations.

337 Sulphur in the Arctic environment (2): results of multi-medium regional mapping
G. KASHULINA, C. REIMANN (Norway)
The regional distribution of sulphur in the terrestrial environment is mainly governed by natural processes.

Number 3 2001

Special Issue: **Terrestrial Carbon — Part I**

Guest Editors: Robert A. Mickler and Steven G. McNulty

351 Preface
R.A. MICKLER, S.G. MCNULTY (USA)

353 Soil carbon dynamics in cropland and rangeland
R. LAL (USA)
Increasing soil carbon sequestration is cost-effective and credible at all levels.

363 Measuring carbon in forests: current status and future challenges
S. BROWN (USA)
Carbon in forests can be measured accurately now, with future measurements relying more on remote sensing and other remote data collecting techniques.

373 Methodology for estimating soil carbon for the forest carbon budget model of the United States, 2001
L.S. HEATH, R.A. BIRDSEY, D.W. WILLIAMS (USA)
Development of soil C pool estimates for the FORCARB model.

381 Carbon storage and sequestration by urban trees in the USA
D.J. NOWAK, D.E. CRANE (USA)
Urban trees currently store 700 million tons of carbon, with an annual sequestration rate of 22.8 million tons.

391 Soil carbon dynamics and potential carbon sequestration by rangelands
G.E. SCHUMAN (USA), H.H. JANZEN (Canada), J.E. HERRICK (USA)
Grazing lands are estimated to contain 10–30% of the world's soil organic carbon.

397 Carbon dioxide fluxes over a grazed prairie and seeded pasture in the Northern Great Plains
A.B. FRANK (USA)
Annual CO_2 flux over temperate grasslands differed, the magnitude depending on dormant season flux.

405 Tillage, crop rotation, and organic amendment effect on changes in soil organic matter
R. RICKMAN, C. DOUGLAS, S. ALBRECHT, J. BERC (USA)
The models CQESTR and RUSLE were used to estimate carbon sequestration in agricultural soils.

413 Sensitivity of carbon sequestration costs to soil carbon rates
J. ANTLE, S. CAPALBO, S. MOONEY, E.T. ELLIOTT, K.H. PAUSTIAN (USA)
Linked economic and biophysical simulation models should be used to assess economic potential for sequestering carbon in agricultural soils.

423 Impact of soil movement on carbon sequestration in agricultural ecosystems
G.W. MCCARTY, J.C. RITCHIE (USA)
Increased sedimentation rates in a wetland ecosystem are associated with higher carbon sequestration rates.

431 National-scale estimation of changes in soil carbon stocks on agricultural lands
M.D. EVE, M. SPEROW, K. PAUSTIAN, R.F. FOLLETT (USA)
Changes in land use and management have resulted in a net gain of 21.2 MMT (million metric tons) of carbon per year in US agricultural soils between 1982 and 1997.

439 Net carbon flux from agricultural ecosystems: methodology for full carbon cycle analyses
T.O. WEST, G. MARLAND (USA)
A methodology was used for analysis that accounts for changes in carbon sequestration, carbon emissions, and net carbon flux with time.

445 Spatial heterogeneity of aggregate stability and soil carbon in semi-arid rangeland
S.B. BIRD, J.E. HERRICK, M.M. WANDER, S.F. WRIGHT (USA)
The authors argue that multi-scale soil heterogeneity must be considered when measuring and managing carbon sequestration.

457 Influence of livestock grazing on C sequestration in semi-arid mixed-grass and short-grass rangelands
J.D. REEDER, G.E. SCHUMAN (USA)
Grazing resulted in higher soil carbon due to more rapid annual shoot turnover and changes in species composition.

Supplement 116 Number 1 2001

Special Issue Supplement to 116/3: **Terrestrial Carbon — Part II**

Guest Editors: Robert A. Mickler and Steven G. McNulty

S1 A carbon balance method for paper and wood products
W.A. CÔTÉ, R.J. YOUNG, K.B. RISSE, A.F. COSTANZA, J.P. TONELLI, C. LENOCKER (USA)
Forest products industries obtaining their raw material from sustainable forest management can achieve a net positive carbon balance over the product life cycle.

S7 Regional estimation of current and future forest biomass
R.A. MICKLER, T.S. EARNHARDT, J.A. MOORE (USA)
Forest net primary production at the regional scale from the PnET model show areas of high and low forest carbon storage potential and their spatial relationship to other landscape features.

S17 Hurricane impacts on US forest carbon sequestration
S.G. McNULTY (USA)
A single hurricane can convert 10% of the total annual carbon storage for the US into dead and downed forest biomass.

S25 Estimating down deadwood from FIA forest inventory variables in Maine
D.C. CHOJNACKY, L.S. HEATH (USA)
The down deadwood component of forest carbon can be modeled at the regional scale as a function of stand size class, basal area of dead and cut trees.

S31 Productivity responses of *Acer rubrum* and *Taxodium distichum* seedlings to elevated CO_2 and flooding
C.D. VANN, J.P. MEGONIGAL (USA)
The growth enhancing effects of elevated atmospheric carbon dioxide were not observed in young seedlings of two tree species in flooded soils due to soil oxygen limitations.

S37 Modeling above-canopy CO_2 flux and evapotranspiration in wheat
P.J. VAUGHAN, D.L. SUAREZ (USA)
Above-ground water vapor flux and carbon dioxide flux were measured and modeled for an Ameriflux wheat field research site.

S45 Anthropogenic and natural CO_2 emission sources in an arid urban environment
B. KOERNER, J. KLOPATEK (USA)
Human and automobile sources produced more than 80% of the carbon dioxide inputs into an urban landscape in a desert environment.

S53 Particulate and non-particulate fractions of soil organic carbon under pastures in the Southern Piedmont USA
A.J. FRANZLUEBBERS, J.A. STUEDEMANN (USA)
Pasture management can improve soil quality and soil carbon sequestration, especially in the particulate organic carbon pools.

S63 Soil carbon protection in podocarp/hardwood forest, and effects of conversion to pasture and exotic pine forest
P.N. BEETS, G.R. OLIVER, P.W. CLINTON (New Zealand)
The effects of land use change on mineral soil carbon were characterized for New Zealand and soil carbon was found to be significantly related to soil pH.

S75 Simulated effects of dryland cropping intensification on soil organic matter and greenhouse gas exchanges using the DAYCENT ecosystem model
S. DEL GROSSO, D. OJIMA, W. PARTON, A. MOSIER, G. PETERSON (USA), D. SCHIMEL (Germany)
The DAYCENT model predicts dryland agricultural soils will be net sources of greenhouse gases but the magnitude can be reduced through management practices.

S85 Influence of seedling roots, environmental factors and soil characteristics on soil CO_2 efflux rates in a 2-year-old loblolly pine (*Pinus taeda* L.) plantation in the Virginia Piedmont
R.E. PANGLE, J. SEILER (USA)
Soil carbon dioxide efflux rates were unchanged following early rotation fertilization while increasing stand growth and carbon sequestration.

S97 Changes in soil quality and below-ground carbon storage with conversion of traditional agricultural crop lands to bioenergy crop production
V.R. TOLBERT, D.E. TODD JR., L.K. MANN, C.M. JAWDY, D.A. MAYS, R. MALIK (USA),
W. BANDARANAYAKE (Canada), A. HOUSTON, D. TYLER, D.E. PETTRY (USA)
The conversion of agricultural lands to bioenergy crops show the potential for increasing long-term soil carbon sequestration and improved soil quality.

S107 Soil carbon pools and fluxes in urban ecosystems
R. POUYAT, P. GROFFMAN, I. YESILONIS, L. HERNANDEZ (USA)
Soil organic carbon pools are directly and indirectly affected by urban landuse conversions and these changes can be observed in adjacent undisturbed forests.

S119 Shade trees reduce building energy use and CO_2 emissions from power plants
H. AKBARI (USA)
Urban tree planting can account for a 25% reduction in net cooling and heating energy usage in urban landscapes.

S127 Spatial variability of soil organic carbon in grasslands: implications for detecting change at different scales
R.T. CONANT, K. PAUSTIAN (USA)
The efficacy of soil sample collection techniques to characterize grassland soil carbon was found to precisely estimate soil carbon changes at the local to national scales.

S137 Effect of soil compaction on organic carbon amounts and distribution, South-Central Iowa
E. BREVIK, T. FENTON, L. MORAN (USA)
The effects of 150 years soil compaction were to alter soil carbon pools by limiting new addditions of organic matter to the soil and limiting vegetative production.

S143 Gaseous carbon dioxide and methane, as well as dissolved organic carbon losses from a small temperate wetland under a changing climate
T.A. CLAIR, P. ARP, T.R. MOORE, M. DALVA, F.-R. MENG (Canada)
The export of dissolved organic carbon from wetlands is projected to double by 2050 under climate change scenarios and increases in atmospheric carbon dioxide

S149 Aspect differences in above- and belowground carbon allocation: a Montana case-study
D. COBLE, J. MARSHALL (USA)
Aboveground net primary production and belowground gross primary production were found to be greater on the north aspect for all leaf area indices in young forest stands in western Montana.

S157 Carbon storage in northeast China as estimated from vegetation and soil inventories
S. WANG, C. ZHOU, J. LIU, H. TIAN, K. LI (People's Republic of China), X. YANG (Japan)
Carbon stocks in soil and vegetation in China are 2.81 PgC and 26.43 PgC respectively with the eastern and northern regions showing the highest carbon storage potential.

S167 Monitoring land-use change effects on soil carbon in New Zealand: quantifying baseline soil carbon stocks
N.A. SCOTT (USA), K.R. TATE, D.J. GILTRAP (New Zealand), C. TATTERSALL SMITH (USA), R.H. WILDE, P.F.J. NEWSOME, M.R. DAVIS (New Zealand)
The national soil carbon stock for New Zealand is estimated at 4,192 MtC using a country-specific land use, climate, and soil carbon system proposed by the IPCC.

S187 Carbon accounting model for forests in Australia
C.L. BRACK, G.P. RICHARDS (Australia)
Carbon sequestration and emissions from Australian forest stands were estimated using the CAMFor model.

S195 Pollution mitigation and carbon sequestration by an urban forest
C.L. BRACK (Australia)
The DISMUT model is a decision support system that has been used to assess Australian urban forest potential for carbon sequestration, pollution mitigation, and energy reduction.

S201 Effects of forest management on soil carbon: results of some long-term resampling studies
D.W. JOHNSON, J.D. KNOEPP, W.T. SWANK, J. SHAN, L.A. MORRIS, D.H. VANLEAR, P.R. KAPELUCK (USA)
The effect of forest harvest treatments on regeneration growth and biomass are likely to cause long-term differences in ecosystem carbon.

S209 Development of a protocol for monitoring status and trends in forest soil carbon at a national level
C.J. PALMER, W.D. SMITH, B.L. CONKLING (USA)
A protocol for monitoring near surface changes in forest soil carbon with bulk density core samplers was evaluated and has been recommended for national implementation.

S221 Using forest health monitoring data to integrate above and below ground carbon information
B.L. CONKLING, C.M. HOOVER, W.D. SMITH, C.J. PALMER (USA)
Field monitoring techniques for national scale characterization of forest carbon budgets have demonstrated the ability to detect a 20% change in total soil carbon and carbon content over 10 years.

S233 Forest inventory and analysis: a national inventory and monitoring program
W.B. SMITH (USA)
The USDA Forest Service Forest Inventory and Analysis program is implementing an integrated approach to monitor and assess the long-term status and trends in the health of U.S. forests.

S243 Local-scale fluxes of carbon dioxide in urban environments: methodological challenges and results from Chicago
C.S.B. GRIMMOND, T.S. KING, F.D. CROPLEY, D.J. NOWAK, C. SOUCH (USA)
The direct measurement and modeling of surface to atmosphere exchanges of carbon dioxide in urban landscapes are lacking to date but will be characterized in the future with continuous eddy covariance measurements.

S255 Use of pyrolysis molecular beam mass spectrometry (py-MBMS) to characterize forest soil carbon: method and preliminary results
K.A. MAGRINI, R.J. EVANS, C.M. HOOVER, C.C. ELAM, M.F. DAVIS (USA)
Changes in soil organic matter chemical composition were successfully characterized using applied pyrolosis molecular beam mass spectrometry on whole soil samples.

S269 Soil carbon content and character in an old-growth forest in northwestern Pennsylvania: a case study introducing pyrolysis molecular beam mass spectrometry (py-MBMS)
C.M. HOOVER, K.A. MAGRINI, R.J. EVANS (USA)
Soil carbon stocks in a chronosequence of forest stands arising from natural disturbance were found to increase with increasing time since disturbance.

S277 The potential of diffuse reflectance spectroscopy for the determination of carbon inventories in soils
J. REEVES III, G. McCARTY (USA), T. MIMMO (Italy)
Near- and mid-infrared reflectance spectrometry techniques can accurately determine organic carbon in soils.

I Contents of Volume 116